

DEHN'S LEMMA FOR IMMERSED LOOPS

MICHAEL FREEDMAN AND MARTIN SCHARLEMANN

Let M be a 3-manifold with boundary ∂M , and $\delta : S^1 \looparrowright \partial M$ be a generic immersion, where $S^1 := [0, 2\pi]/0 \equiv 2\pi$. By generic, we mean that δ has only simple crossings. We say that δ' is δ “displaced by a height function f ” if $\delta'(\theta) = (\delta(\theta), f(\theta))$, where $f : S^1 \rightarrow (0, \varepsilon)$ is a Morse function and $[0, \varepsilon]$ is a normal collar coordinate on ∂M into M . We call a simple closed curve (scc) $\alpha \subset \text{int}(M)$ unknotted iff α bounds an embedded disk $\bar{\alpha} : D^2 \rightarrow \text{int}(M)$.

Theorem 1. *Let $\delta : S^1 \looparrowright \partial M$ be a generic immersed loop so that the composition into M is null homotopic. There is a height function $f : S^1 \rightarrow (0, \varepsilon)$ so that $\delta' = (\delta, \varepsilon) : S^1 \hookrightarrow \text{int}(M)$ is unknotted.*

The theorem readily implies two familiar facts:


- (1) Dehn’s Lemma: For any scc $\delta \subset \partial M$ which is null homotopic in M there is a properly embedded disk $(D, \partial D) \subset (M, \partial M)$ with ∂D parameterizing δ .
- (2) Given any knot diagram there is always a way to rechoose the crossings to produce an unknot. (This is the case when $(M, \partial M) \cong (B^3, \partial B^3)$ is a 3-ball.)

Regarding 1: A short argument connects the special case of the theorem where δ is one-to-one to Dehn’s Lemma. Let C be the annular collar joining δ to δ' and D a disk with boundary δ' . If C and D are in general position there is a local integral linking number l between the normals of δ' in C vs D . Perturbing D near δ' allows l to be made zero. A further perturbation of D near δ' ensures that C and D intersect only in sccs contains in $\text{int}(C)$ and $\text{int}(D)$. Let $\sigma \subset D$ be an innermost circle of intersection bounding a subdisk $\bar{\sigma} \in D$. σ may be paired with either an essential or inessential scc in C . Perform disk exchanges to modify C until either $C \cup D$ is an embedded disk or some innermost σ is paired to an essential scc in C . In this case a final cut and glue operation yields an embedded disk with boundary δ .


Regarding 2: We should note a subtlety. The height function produced in the theorem may, in general, be more complicated than the familiar height function which solves the knot diagram problem (see Figure 1b). For knot diagrams the unknotting function may be taken to be any function with a unique local maximum and unique local minimum.

Actually Theorem 1 is a corollary of a stronger Theorem 0, better adapted to the required induction.

Theorem 0. *Let $\delta : S^1 \looparrowright \partial M$ be a generic immersed loop with base point $*$ ($*$ is assumed disjoint from multiple points of δ) whose composition into M is null homotopic. There is a height function $f : S^1 \rightarrow [0, \varepsilon]$, $f(*) = 0$, $f(S^1 \setminus *) \subset (0, \varepsilon)$, so that $\delta' = \{(\delta(\theta), f(\theta))\}$ bounds an embedded disk $\Delta \subset M$ with $\Delta \cap \partial M = *$. We say δ' collapses to $*$, and call Δ a “lollypop” for δ' .*



(A) This example of a loop on the boundary of a genus 2 handlebody shows that the height function f , constructed in the proof of Theorem 0, cannot always have a unique local maximum, the familiar form for unknotting knot diagrams on the 2-sphere. The unknotting function f , relative to the indicated base point $*$ with $f(*) = 0$, must have at least two local maxima.



(B) For this absolute example, the unknotting function f produced by Theorem 1 must have at least m local minima (and maxima)

FIGURE 1

Proof. No methods post dating Papakyriakopoulos [1957]¹ are required. He would have found this proof rather easy to understand and perhaps to generate. First we build a tower.

δ is the pointed, immersed loop, D the immersed null homotopy bounding it, and N the regular neighborhood of D in M . Subscript will indicate height in the tower.

To build the tower we should ask about the Z_2 Betti number $b_1(N) := \text{rank}(H_1(N; Z_2))$. If $b_1(N) = 0$, there is no tower. For homological reasons ∂N is a disjoint union of 2-spheres and δ is contained in one of these: $\delta \looparrowright S \subset \partial N$.

If $b_1(N) > 0$ choose a 2-fold cover $\tilde{N}_0 \rightarrow N_0 := N$ and choose a lift $l_1 : D \looparrowright \tilde{N}_0$ and let $N_1 = \text{neib}(l_1(D)) \subset \tilde{N}_0$. If $b_1(N_1) = 0$ then N_1 is the top of the tower. If $b_1(N_1) > 0$ continue and find a 2-fold cover $\tilde{N}_1 \rightarrow N_1$, and lift $l_2 : D \looparrowright \tilde{N}_1$ and set $N_2 = \text{neib}(l_2(D)) \subset \tilde{N}_1$. Again if $b_1(N_2) = 0$ then N_2 is the top of the tower. If not, proceed to construct $l_3 \dots l_n$ and $N_3 \dots N_n$ until $b_1(N_n) = 0$. A simple complexity argument, where the complexity can be the number of simplices identified by l_k for a fixed triangulation of D making all l_k simplicial, shows that the tower is indeed finite.

¹Papakyriakopoulos, C. D. (1957b), "On Dehn's Lemma and the Asphericity of Knots", Ann. Math., 66 (1): 126.




FIGURE 2. The tower

Observation: $\delta_n = l_n(\partial D)$ can be unknotted by a suitable resolution of its crossings (i.e. a normal function f as in Theorem 1). As in Theorem 0, given for any base point $*$ (chosen away from crossings) we can resolve crossings $\delta_n \rightarrow \delta'_n$ and produce an embedded disk Δ with $\partial\Delta = \delta'_n$ and $\Delta \cap \partial N_n = *$.

Proof. One might expect to choose $\delta'_n = \{\delta_n(\theta), f(\theta)\}$, $f(*) = 0$, $f(\theta \neq *) > 0$, where f has a unique local maximum and a unique local minimum. However, this solution does not, in general, push down the tower. We prefer to give a second solution. There are two cases. If l_n is one-to-one it bounds a hemisphere $H \subset S \subset \partial N_n$. Pushing H normally toward interior N_n gives a disk Δ bounding δ'_n .

Now assume δ_n is not one-to-one. Let $\alpha \subset \delta_n$, not containing $*$, be a subarc so that $l_n(\alpha)$ is a scc $\subset \partial N$. Begin to resolve the crossings of δ by following this rule: $l_n(\alpha)$ lies above $l_n(\beta)$, where β is the complementary arc, $\delta = \alpha \cup \beta$ and $\alpha \cap \beta = \partial\alpha = \partial\beta$. Also the two endpoints of α are at a crossing: resolve this crossing arbitrarily.




FIGURE 3

Now an isotopy across a 2-cell C bounding α effectively erases the loop α . If the simplified diagram is a scc we may continue the unknotting using a hemisphere cell in S making no further crossing choices. If the simplified diagram is still singular choose another arc $\alpha' \subset \beta$ whose image is again a scc not containing $*$ and proceed as before. Continuing in this way, guided by a sequence of embedded 2-cells, say $\{E_1, \dots, E_j\}$, all crossings are eventually assigned so that the diagram resolution is unknotted with $*$ remaining in the final 2-cell E_j . \square

This solution will now be pushed down the tower

Dissection of disks with double arcs: Let E be a properly embedded disk $(E, \partial E) \subset (M^3, \partial M)$ in a 3-manifold and $\pi : (M, \partial M) \rightarrow (P, \partial P)$ a 2-fold cover with covering translation t . Assume E and tE are in generic position and meet only in double arcs and double loops. Double loops are easily removed by an innermost circle argument and will not be discussed further. E is assumed to have a base point $* \subset \partial E$. Call $\pi(E) = F$ and $\pi(*) = *$ in an abuse of notation.

We now describe how to form an ordered list of embedded disks $F_1, \dots, F_k \subset (P, \partial)$ from pieces (some used several times) of F . Constructing $\{F_1, \dots, F_k\}$ will dictate “crossing choices” for $\partial F := \gamma$, which yield γ' , bounding a lollipop. Each F_i has base point $*_i$ with $*_k = * \subset \partial F$. Furthermore, once $\{F_1, \dots, F_k\}$ are constructed we will see that they constitute the instructions for collapsing γ' to $*$.

Let β_1 be an outermost arc cutting off an outermost disk $D_1 \subset F$ not containing $*$. Let β'_1 be the partner arc of β_1 and D'_1 be the subdisk of F cut off by β'_1 which also does not contain $*$. There are two cases:





FIGURE 4

In both cases $D_1 \cup D'_1$ can be perturbed into a proper map of a disk F_0 . There is a residual general position proper map with only double arc singularities of a disk formed from $(F \setminus D'_1) \cup D_1$ in both cases, temporarily label this map F_1 . Because D_1 is outermost both F_0 and F_1 have fewer double arcs than F .

We need to discuss certain crossing choices (resolutions) and base point choices. All crossing points on $\partial D'_1$ with other segments of ∂F are deemed overcrossings (in both cases). F_0 is provided a base point $*_0$ in its copy of D_1 and F_1 retains $*_1 := *$ as its base point.

Now unless $F_0(F_1)$ is embedded find an outermost arc cutting off an outermost disk D , with D' disjoint from $*_0(*_1)$ and sharing a double arc with D , not containing $*_0(*_1)$ and as above dissect:



to obtain general position proper pointed maps F_{00}, F_{01}, F_{10} , and F_{11} of disks with only double arc singularities. We call such maps good maps. In each case follow the preceding rule for resolving certain double points on the boundary (crossings) as overcrossings. Specifically, all crossings of $\partial D'$ with other segments of ∂F are deemed overcrossings.

Continuing in this way a diatic tree of good maps is obtained. The leaves of this tree are called great maps as they have the additional property of being embeddings. (No disjointness has been assumed or constructed.) The leaves are now linearly ordered by the base 2 numerical value of their subscripts considered as decimals. By the time we reach the leaves all crossings have been resolved. These great disks, monotonically reindexed becomes the ordered list $\{F_1, \dots, F_k\}$, build from pieces of F_1 , which we sought. We call the result a great sequence (of disks) guiding a collapse of $(\partial F)'$ to $*$. The simple expedient of successively declaring $\partial D'$ to over-cross other segments of ∂F has given us the well-defined crossing choices defining $(\partial F)'$.




FIGURE 5. Sample tree with great maps as leaves

We need some terminology.

Definition 2. *The tree, as constructed above, of good disks terminating in great disks is a dissection of F . The same tree, according to our convention, determines crossing choices γ' for $\gamma = \partial F$. The great sequence $\{F_1, \dots, F_k\}$ (the leaves of the dissection) are said to guide a sequence of isotopes $I_k \circ \dots \circ I_1 := I$ with I_i supported near F_i , $1 \leq i \leq k$, so that I collapses γ' to its base point.*

Example: We illustrate (Figure 6) dissection, crossing resolution, and the collapsing isotopy with an example. Our crossing convention implies that middle sheet of the three sheets of D_1 in $F_0 \amalg F_1$ should be compressed slightly into interior (M) near $\partial \beta_1$ when interpreting the disk F_j , $1 \leq j \leq k$, as guiding the isotopes I_j .

We now finish the proof of Theorem 0.

At the top of the tower the collapse of $((I_n(\partial D))', *)$ is guided by an initial great sequence (of properly embedded 2-cells) $\{E_1, \dots, E_j\}$, where the base point of each E_1, \dots, E_j is taken to be the self intersection point of its boundary, and with the original $* \subset E_j$ serving as the base point for the final E_j .




FIGURE 6

Now consider each of these cells E_i , $1 \leq i \leq j$, and its singular image F_i under the 2-fold cover $\pi_{n-1} : \tilde{N}_{n-1} \rightarrow N_{n-1}$. Again we may remove double loops and only need consider double arcs. As above, dissect F_i into a sequence of great disks $\{F_{i,1}, \dots, F_{i,k_i}\}$ and replace each E_i with that ordered sublist to obtain a great sequence:

$$\{F_{1,1}, \dots, F_{1,k_1}, \dots, F_{j,1}, \dots, F_{j,k_j}\}$$

which guides the collapse of $(l_{n-1}(\partial D))'$ to $*$.

Now proceed all the way down the tower. At height m by induction assume a great sequence $\{G_1, \dots, G_p\}$ for $(l_m(\partial D))'$. Projecting by π_{m-1} , each G_i , $1 \leq i \leq p$, to N_{m-1} we construct, as above, a great sequence for $(\pi_{m-1}(\partial G_i))'$. The great sequence for $(l_{m-1}(\partial D))'$ is obtained by concatenating these p great sequences. \square

MICHAEL FREEDMAN, MICROSOFT RESEARCH, STATION Q, AND DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, SANTA BARBARA, CA 93106,

E-mail address: mfreedman@math.ucsb.edu

MARTIN SCHARLESMANN, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA BARBARA, SANTA BARBARA, CA 93106,

E-mail address: mgscharl@math.ucsb.edu